Noname manuscript No.
(will be inserted by the editor)

Symmetric identities of higher-order degenerate Euler polynomials

Dae San Kim · Taekyun Kim

arXiv:1704.04025v1 [math.NT] 13 Apr 2017

Received: date / Accepted: date

Abstract The purpose of this paper is to give some symmetric identities of higher-order degenerate Euler polynomials derived from the symmetric properties of the multivariate p-adic fermionic integrals on $\mathbb{Z}_p$.

Keywords Symmetry · Higher-order degenerate Euler polynomial · Identity

1 Introduction

Let p be a fixed prime such that $p \equiv 1 \pmod 2$. Throughout this paper, $\mathbb{Z}_p$, $\mathbb{Q}_p$, and $\mathbb{C}_p$ will denote the ring of p-adic integers, the field of p-adic numbers and the completion of the algebraic closure of $\mathbb{Q}_p$, respectively. Let ν_p be the normalized exponential valuation of $\mathbb{C}_p$ with $|p|_p = p^{-\nu_p(p)} = p^{-1}$.

Let $f(x)$ be a continuous function on $\mathbb{Z}_p$. Then p-adic fermionic integral on $\mathbb{Z}_p$ is defined by Kim as

$$\begin{aligned} \int_{\mathbb{Z}_p} f(x)\, d\mu_{-1}(x) &= \lim_{N\to\infty} \sum_{x=0}^{p^N-1} f(x)\, \mu_{-1}\left(x + p^N\mathbb{Z}_p\right) \\ &= \lim_{N\to\infty} \sum_{x=0}^{p^N-1} f(x)\,(-1)^x, \quad (\text{see } [8]). \end{aligned} \tag{1}$$

Dae San Kim
Department of Mathematics, Sogang University, Seoul 121-742, Republic of Korea
Tel.: +82-2-7058871
Fax: +82-2-7058871
E-mail: dskim@sogang.ac.kr

Taekyun Kim
Department of Mathematics, Kwangwoon University, Seoul 139-701, Republic of Korea
E-mail: tkkim@kw.ac.kr

From (1), we note that

$$\int_{\mathbb{Z}_p} f(x+1)\, d\mu_{-1}(x) + \int_{\mathbb{Z}_p} f(x)\, d\mu_{-1}(x) = 2f(0), \tag{2}$$

and

$$\begin{aligned}&\int_{\mathbb{Z}_p} f(x+n)\, d\mu_{-1}(x) + (-1)^{n-1} \int_{\mathbb{Z}_p} f(x)\, d\mu_{-1}(x)\\ &= 2\sum_{l=0}^{n-1} (-1)^{n-1-l} f(l), \quad (\text{see } [4]).\end{aligned} \tag{3}$$

As is well known, the Euler polynomials are defined by the generating function

$$\frac{2}{e^t+1} e^{xt} = \sum_{n=0}^{\infty} E_n(x) \frac{t^n}{n!}, \quad (\text{see } [4,8]). \tag{4}$$

When $x = 0$, $E_n = E_n(0)$ are called the Euler numbers.

For $r \in \mathbb{N}$, the higher-order Euler polynomials are given by

$$\left(\frac{2}{e^t+1}\right)^r e^{xt} = \sum_{n=0}^{\infty} E_n^{(r)}(x) \frac{t^n}{n!}, \quad (\text{see } [1,2,3,4,5,6,7,9,10]) \tag{5}$$

In particular, $x = 0$, $E_n^{(r)} = E_n^{(r)}(0)$ are called the Euler numbers of order r.

From (2), we can easily derive the following equation:

$$\begin{aligned}&\int_{\mathbb{Z}_p} \cdots \int_{\mathbb{Z}_p} e^{(x_1+\cdots+x_r+x)t} d\mu_{-1}(x_1) \cdots d\mu_{-1}(x_r)\\ &= \left(\frac{2}{e^t+1}\right)^r e^{xt}\\ &= \sum_{n=0}^{\infty} E_n^{(r)}(x) \frac{t^n}{n!}.\end{aligned} \tag{6}$$

Thus, by (6), we get

$$\int_{\mathbb{Z}_p} \cdots \int_{\mathbb{Z}_p} (x_1 + \cdots + x_r + x)^n\, d\mu_{-1}(x_1) \cdots d\mu_{-1}(x_r) = E_n^{(r)}(x), \quad (n \geq 0). \tag{7}$$

Carlitz introduced the degenerate Euler polynomials given by the generating function

$$\left(\frac{2}{(1+\lambda t)^{\frac{1}{\lambda}}+1}\right)(1+\lambda t)^{\frac{x}{\lambda}} = \sum_{n=0}^{\infty} \mathcal{E}_n(x \mid \lambda) \frac{t^n}{n!}, \quad (\text{see } [5]). \tag{8}$$

When $x = 0$, $\mathcal{E}_n(0 \mid \lambda) = \mathcal{E}_n(\lambda)$ are the degenerate Euler numbers. Note that $\lim_{\lambda \to 0} \mathcal{E}_{n,\lambda}(x \mid \lambda) = E_n(x)$.

For $r \in \mathbb{N}$, the higher-order degenerate Euler polynomials are also given by the generating function

$$\left(\frac{2}{(1+\lambda t)^{\frac{1}{\lambda}}+1}\right)^r (1+\lambda t)^{\frac{x}{\lambda}} = \sum_{n=0}^{\infty} \mathcal{E}_n^{(r)}(x \mid \lambda) \frac{t^n}{n!}. \tag{9}$$

When $x = 0$, $\mathcal{E}_n^{(r)}(0 \mid \lambda) = \mathcal{E}_n^{(r)}(\lambda)$ are called the higher-order degenerate Euler numbers (see [2]).

In [8], Kim and Kim showed that the degenerate Euler polynomials can be represented by a p-adic integral on $\mathbb{Z}_p$. Recently, several researchers have studied the symmetric identities of higher-order Euler polynomials derived from the symmetric properties of p-adic integrals on $\mathbb{Z}_p$(see [1,2,3,4,5,6,7,8, 9,10,11,12,13,14,15,16,17,18,19]).

In this paper, we investigate some properties of symmetry for the multivariate p-adic fermionic integrals on $\mathbb{Z}_p$. From our investigation, we derive some identities of symmetry for the higher-order degenerate Euler polynomials.

2 Identities of symmetry for the higher-order degenerate Euler polynomials

In this section, we assume that $\lambda, t \in \mathbb{C}_p$ with $|\lambda t|_p < p^{-\frac{1}{p-1}}$. From (2), we can derive the following equation:

$$\begin{aligned} &\int_{\mathbb{Z}_p} \cdots \int_{\mathbb{Z}_p} (1+\lambda t)^{\frac{x_1+\cdots+x_r+x}{\lambda}} d\mu_{-1}(x_1) \cdots d\mu_{-1}(x_r) \\ &= \left(\frac{2}{(1+\lambda t)^{\frac{1}{\lambda}}+1}\right)^r (1+\lambda t)^{\frac{x}{\lambda}} \\ &= \sum_{n=0}^{\infty} \mathcal{E}_n^{(r)}(x \mid \lambda) \frac{t^n}{n!}. \end{aligned} \tag{10}$$

Thus, by (10), we get, for $n \geq 0$,

$$\int_{\mathbb{Z}_p} \cdots \int_{\mathbb{Z}_p} (x_1 + \cdots + x_r + x \mid \lambda)_n \, d\mu_{-1}(x_1) \cdots d\mu_{-1}(x_r) = \mathcal{E}_n^{(r)}(x \mid \lambda), \tag{11}$$

where

$$(x \mid \lambda)_n = x(x-\lambda) \cdots (x-\lambda(n-1)), \quad (n \geq 0). \tag{12}$$

From (12), we note that

$$(x \mid \lambda)_n = \sum_{l=0}^{n} S_1(n,l) \lambda^{n-l} x^l, \quad (n \geq 0), \tag{13}$$

where $S_1(n,l)$ is the Stirling number of the first kind.

From (3), we have, for $m \geq 0$,

$$2\sum_{l=0}^{n-1}(-1)^{n-1-l}(l \mid \lambda)_m \tag{14}$$
$$= \int_{\mathbb{Z}_p}(x+n \mid \lambda)_m \, d\mu_{-1}(x) + (-1)^{n-1}\int_{\mathbb{Z}_p}(x \mid \lambda)_m \, d\mu_{-1}(x).$$

For $n \in \mathbb{N}$ with $n \equiv 1 \pmod 2$, we get

$$2\sum_{l=0}^{n-1}(-1)^l (l \mid \lambda)_m = \mathcal{E}_m(n \mid \lambda) + \mathcal{E}_m(\lambda), \quad (m \geq 0). \tag{15}$$

Let us define $\tilde{S}_k(n \mid \lambda)$ as follows:

$$\tilde{S}_k(n \mid \lambda) = \sum_{l=0}^{n}(-1)^l (l \mid \lambda)_k, \quad (k, n \geq 0). \tag{16}$$

Then, we note that

$$\lim_{\lambda \to 0}\tilde{S}_k(n \mid \lambda) = \sum_{l=0}^{n}(-1)^l l^k = \tilde{S}_k(n). \tag{17}$$

Let $n \in \mathbb{N}$ with $n \equiv 1 \pmod 2$. Then we get

$$\begin{aligned} &\frac{2\int_{\mathbb{Z}_p}(1+\lambda t)^{\frac{x}{\lambda}} d\mu_{-1}(x)}{\int_{\mathbb{Z}_p}(1+\lambda t)^{\frac{nx}{\lambda}} d\mu_{-1}(x)} \\ &= \int_{\mathbb{Z}_p}(1+\lambda t)^{\frac{n+x}{\lambda}} d\mu_{-1}(x) + \int_{\mathbb{Z}_p}(1+\lambda t)^{\frac{x}{\lambda}} d\mu_{-1}(x) \\ &= 2\sum_{l=0}^{n-1}(-1)^l (1+\lambda t)^{\frac{l}{\lambda}} \\ &= 2\sum_{k=0}^{\infty}\tilde{S}_k(n-1 \mid \lambda)\frac{t^k}{k!}. \end{aligned} \tag{18}$$

Let $w_1, w_2 \in \mathbb{N}$, with $w_1 \equiv 1 \pmod 2$, $w_2 \equiv 1 \pmod 2$. For $m \in \mathbb{N}$, we define

$$\begin{aligned} &K^{(m)}(w_1, w_2 \mid \lambda) \\ &= \left(\frac{2}{(1+\lambda t)^{\frac{w_1}{\lambda}}+1}\right)^m (1+\lambda t)^{\frac{w_1 w_2 x}{\lambda}}\left((1+\lambda t)^{\frac{w_1 w_2}{\lambda}}+1\right) \\ &\quad \times \left(\frac{2}{(1+\lambda t)^{\frac{w_2}{\lambda}}+1}\right)^m \frac{1}{2}(1+\lambda t)^{\frac{w_1 w_2}{\lambda}y}. \end{aligned} \tag{19}$$

From (3), we note that

$$K^{(m)}(w_1,w_2\mid\lambda)=\frac{\int_{\mathbb{Z}_p^m}(1+\lambda t)^{\frac{w_1}{\lambda}(x_1+\cdots+x_m+w_2x)}\,d\mu_{-1}(x_1)\cdots d\mu_{-1}(x_m)}{\int_{\mathbb{Z}_p}(1+\lambda t)^{\frac{w_1w_2}{\lambda}x}\,d\mu_{-1}(x)} \tag{20}$$
$$\times\int_{\mathbb{Z}_p^m}(1+\lambda t)^{\frac{w_2}{\lambda}(x_1+\cdots+x_m+w_1y)}\,d\mu_{-1}(x_1)\cdots d\mu_{-1}(x_m),$$

where

$$\int_{\mathbb{Z}_p^m}f(x_1,\ldots,x_m)\,d\mu_{-1}(x_1)\cdots d\mu_{-1}(x_m) \tag{21}$$
$$=\int_{\mathbb{Z}_p}\cdots\int_{\mathbb{Z}_p}f(x_1,\ldots,x_m)\,d\mu_{-1}(x_1)\cdots d\mu_{-1}(x_m).$$

It is easy to see that $K^{(m)}(w_1,w_2\mid\lambda)$ is symmetric in w_1 and w_2. Now, we observe that

$$K^{(m)}(w_1,w_2\mid\lambda) \tag{22}$$
$$=(1+\lambda t)^{\frac{w_1w_2}{\lambda}x}\int_{\mathbb{Z}_p^m}(1+\lambda t)^{\frac{w_1}{\lambda}(x_1+\cdots+x_m)}\,d\mu_{-1}(x_1)\cdots d\mu_{-1}(x_m)$$
$$\times\frac{\int_{\mathbb{Z}_p}(1+\lambda t)^{\frac{w_2}{\lambda}x_m}\,d\mu_{-1}(x_m)}{\int_{\mathbb{Z}_p}(1+\lambda t)^{\frac{1}{\lambda}w_1w_2x}\,d\mu_{-1}(x)}$$
$$\times(1+\lambda t)^{\frac{1}{\lambda}w_1w_2y}\int_{\mathbb{Z}_p^{m-1}}(1+\lambda t)^{\frac{w_2}{\lambda}(x_1+\cdots+x_{m-1})}\,d\mu_{-1}(x_1)\cdots d\mu_{-1}(x_{m-1}).$$

By (10), we get

$$(1+\lambda t)^{\frac{w_1w_2}{\lambda}x}\int_{\mathbb{Z}_p^m}(1+\lambda t)^{\frac{w_1}{\lambda}(x_1+\cdots+x_m)}\,d\mu_{-1}(x_1)\cdots d\mu_{-1}(x_m) \tag{23}$$
$$=\left(\frac{2}{(1+\lambda t)^{\frac{w_1}{\lambda}}+1}\right)^m(1+\lambda t)^{\frac{w_1w_2}{\lambda}x}$$
$$=\sum_{n=0}^{\infty}\mathcal{E}_n^{(m)}\left(w_2x\,\middle|\,\frac{\lambda}{w_1}\right)w_1^n\frac{t^n}{n!}.$$

From (18), (22) and (23), we have

$$K^{(m)}(w_1,w_{2,}\mid\lambda) \tag{24}$$
$$=\sum_{l=0}^{\infty}\mathcal{E}_l^{(m)}\left(w_2x\,\middle|\,\frac{\lambda}{w_1}\right)w_1^l\frac{t^l}{l!}\sum_{k=0}^{\infty}\tilde{S}_k\left(w_1-1\,\middle|\,\frac{\lambda}{w_2}\right)\frac{w_2^k}{k!}t^k$$
$$\times\sum_{i=0}^{\infty}\mathcal{E}_i^{(m-1)}\left(w_1y\,\middle|\,\frac{\lambda}{w_2}\right)\frac{w_2^i}{i!}t^i$$

$$= \sum_{l=0}^{\infty} \mathcal{E}_l^{(m)} \left(w_2 x \left| \frac{\lambda}{w_1} \right. \right) w_1^l \frac{t^l}{l!}$$
$$\times \sum_{j=0}^{\infty} \sum_{k=0}^{j} \tilde{S}_k \left(w_1 - 1 \left| \frac{\lambda}{w_2} \right. \right) \binom{j}{k} w_2^k w_2^{j-k} \mathcal{E}_{j-k}^{(m-1)} \left(w_1 y \left| \frac{\lambda}{w_2} \right. \right) \frac{t^j}{j!}$$
$$= \sum_{n=0}^{\infty} \sum_{j=0}^{n} \binom{n}{j} w_2^j w_1^{n-j} \mathcal{E}_{n-j}^{(m)} \left(w_2 x \left| \frac{\lambda}{w_1} \right. \right)$$
$$\times \sum_{k=0}^{j} \tilde{S}_k \left(w_1 - 1 \left| \frac{\lambda}{w_2} \right. \right) \binom{j}{k} \mathcal{E}_{j-k}^{(m-1)} \left(w_1 y \left| \frac{\lambda}{w_2} \right. \right) \frac{t^n}{n!}.$$

On the other hand,

$$K^{(m)}(w_1, w_2 \mid \lambda) \tag{25}$$
$$= \sum_{n=0}^{\infty} \sum_{j=0}^{n} \binom{n}{j} w_1^j w_2^{n-j} \mathcal{E}_{n-j}^{(m)} \left(w_1 x \left| \frac{\lambda}{w_2} \right. \right)$$
$$\times \sum_{k=0}^{j} \tilde{S}_k \left(w_2 - 1 \left| \frac{\lambda}{w_1} \right. \right) \binom{j}{k} \mathcal{E}_{j-k}^{(m-1)} \left(w_2 y \left| \frac{\lambda}{w_1} \right. \right) \frac{t^n}{n!}.$$

Therefore, by (24) and (25), we obtain the following theorem.

Theorem 1 *For $w_1, w_2 \in \mathbb{N}$, with $w_1 \equiv 1 \pmod 2$, $w_2 \equiv 1 \pmod 2$, $n \geq 0$ and $m \in \mathbb{N}$, we have*

$$\sum_{j=0}^{n} \binom{n}{j} w_2^j w_1^{n-j} \mathcal{E}_{n-j}^{(m)} \left(w_2 x \left| \frac{\lambda}{w_1} \right. \right) \sum_{k=0}^{j} \tilde{S}_k \left(w_1 - 1 \left| \frac{\lambda}{w_2} \right. \right) \binom{j}{k} \mathcal{E}_{j-k}^{(m-1)} \left(w_1 y \left| \frac{\lambda}{w_2} \right. \right)$$
$$= \sum_{j=0}^{n} \binom{n}{j} w_1^j w_2^{n-j} \mathcal{E}_{n-j}^{(m)} \left(w_1 x \left| \frac{\lambda}{w_2} \right. \right) \sum_{k=0}^{j} \tilde{S}_k \left(w_2 - 1 \left| \frac{\lambda}{w_1} \right. \right) \binom{j}{k} \mathcal{E}_{j-k}^{(m-1)} \left(w_2 y \left| \frac{\lambda}{w_1} \right. \right).$$

Let $y = 0$ and $m = 1$ in Theorem 1. Then we have the following theorem.

Theorem 2 *For $n \geq 0$, $w_1, w_2 \in \mathbb{N}$ with $w_1 \equiv 1, w_2 \equiv 1 \pmod 2$, we have*

$$\sum_{j=0}^{n} \binom{n}{j} w_2^j w_1^{n-j} \mathcal{E}_{n-j} \left(w_2 x \left| \frac{\lambda}{w_1} \right. \right) \tilde{S}_j \left(w_1 - 1 \left| \frac{\lambda}{w_2} \right. \right)$$
$$= \sum_{j=0}^{n} \binom{n}{j} w_1^j w_2^{n-j} \mathcal{E}_{n-j} \left(w_1 x \left| \frac{\lambda}{w_2} \right. \right) \tilde{S}_j \left(w_2 - 1 \left| \frac{\lambda}{w_1} \right. \right).$$

In particular, if we take $w_2 = 1$ in Theorem 2, then we obtain the following corollary.

Corollary 3 *For $w_1 \in \mathbb{N}$ with $w_1 \equiv 1 \pmod 2$, $n \geq 0$, we have*

$$\mathcal{E}_n(w_1 x \mid \lambda) = \sum_{j=0}^{n} \binom{n}{j} w_1^{n-j} \mathcal{E}_{n-j} \left(x \left| \frac{\lambda}{w_1} \right. \right) \tilde{S}_j(w_1 - 1 \mid \lambda).$$

From (19), we have

$$\begin{aligned}
&K^{(m)}\left(w_1,w_2 \mid \lambda\right) \qquad (26)\\
&=\left(\frac{2}{\left(1+\lambda t\right)^{\frac{w_1}{\lambda}}+1}\right)^m \left(1+\lambda t\right)^{\frac{w_1w_2}{\lambda}x}\\
&\quad\times\frac{\left(1+\lambda t\right)^{\frac{w_1w_2}{\lambda}}+1}{\left(1+\lambda t\right)^{\frac{w_2}{\lambda}}+1}\left(\frac{2}{\left(1+\lambda t\right)^{\frac{w_2}{\lambda}}+1}\right)^{m-1}\left(1+\lambda t\right)^{\frac{w_1w_2}{\lambda}y}\\
&=\left(\frac{2}{\left(1+\lambda t\right)^{\frac{w_1}{\lambda}}+1}\right)^m \sum_{i=0}^{w_1-1}\left(-1\right)^i\left(1+\lambda t\right)^{\frac{w_1w_2}{\lambda}x+\frac{w_2}{\lambda}i}\\
&\quad\times\left(\frac{2}{\left(1+\lambda t\right)^{\frac{w_2}{\lambda}}+1}\right)^{m-1}\left(1+\lambda t\right)^{\frac{w_1w_2}{\lambda}y}\\
&=\sum_{i=0}^{w_1-1}\left(-1\right)^i\sum_{k=0}^{\infty}\mathcal{E}_k^{(m)}\left(w_2x+\frac{w_2}{w_1}i\middle|\,\frac{\lambda}{w_1}\right)w_1^k\frac{t^k}{k!}\\
&\quad\times\sum_{l=0}^{\infty}\mathcal{E}_l^{(m-1)}\left(w_1y|\,\frac{\lambda}{w_2}\right)w_2^l\frac{t^l}{l!}\\
&=\sum_{n=0}^{\infty}\sum_{k=0}^{n}\binom{n}{k}\sum_{i=0}^{w_1-1}\left(-1\right)^i\mathcal{E}_k^{(m)}\left(w_2x+\frac{w_2}{w_1}i\middle|\,\frac{\lambda}{w_1}\right)w_1^k\\
&\quad\times\mathcal{E}_{n-k}^{(m-1)}\left(w_1y|\,\frac{\lambda}{w_2}\right)w_2^{n-k}\frac{t^n}{n!}.
\end{aligned}$$

On the other hand, by the symmetric properties of $K^{(m)}\left(w_1,w_2 \mid \lambda\right)$ in w_1 and w_2, we get

$$\begin{aligned}
&K^{(m)}\left(w_1,w_2 \mid \lambda\right) \qquad (27)\\
&=\sum_{n=0}^{\infty}\sum_{k=0}^{n}\binom{n}{k}w_2^kw_1^{n-k}\mathcal{E}_{n-k}^{(m-1)}\left(w_2y|\,\frac{\lambda}{w_1}\right)\\
&\quad\times\sum_{i=0}^{w_2-1}\left(-1\right)^i\mathcal{E}_k^{(m)}\left(w_1x+\frac{w_1}{w_2}i\middle|\,\frac{\lambda}{w_2}\right)\frac{t^n}{n!}.
\end{aligned}$$

Therefore, by comparing the coefficients on the both sides of (26) and (27), we obtain the following theorem.

Theorem 4 *For $w_1,w_2\in\mathbb{N}$ with $w_1\equiv 1, w_2\equiv 1 \pmod 2$ and $n\geq 0$ and $m\geq 1$, we have*

$$\begin{aligned}
&\sum_{k=0}^{n}\binom{n}{k}w_1^kw_2^{n-k}\mathcal{E}_{n-k}^{(m-1)}\left(w_1y|\,\frac{\lambda}{w_2}\right)\sum_{i=0}^{w_1-1}\left(-1\right)^i\mathcal{E}_k^{(m)}\left(w_2x+\frac{w_2}{w_1}i\middle|\,\frac{\lambda}{w_1}\right)\\
&=\sum_{k=0}^{n}\binom{n}{k}w_2^kw_1^{n-k}\mathcal{E}_{n-k}^{(m-1)}\left(w_2y|\,\frac{\lambda}{w_1}\right)\sum_{i=0}^{w_2-1}\left(-1\right)^i\mathcal{E}_k^{(m)}\left(w_1x+\frac{w_1}{w_2}i\middle|\,\frac{\lambda}{w_2}\right).
\end{aligned}$$

Let $y = 0$ and $m = 1$ in Theorem 4. Then we have the following corollary.

Corollary 5 *For $w_1, w_2 \in \mathbb{N}$ with $w_1 \equiv 1, w_2 \equiv 1 \pmod 2$, $n \geq 0$, we have*

$$w_1^n \sum_{i=0}^{w_1-1} (-1)^i \mathcal{E}_n \left(w_2 x + \frac{w_2}{w_1} i \middle| \frac{\lambda}{w_1} \right) = w_2^n \sum_{i=0}^{w_2-1} (-1)^i \mathcal{E}_n \left(w_1 x + \frac{w_1}{w_2} i \middle| \frac{\lambda}{w_2} \right).$$

Let us take $w_2 = 1$ in Corollary 5. Then we have

$$w_1^n \sum_{i=0}^{w_1-1} (-1)^i \mathcal{E}_n \left(x + \frac{1}{w_1} i \middle| \frac{\lambda}{w_1} \right) = \mathcal{E}_n \left(w_1 x \mid \lambda \right).$$

References

1. Araci, S., Acikgoz, M.: A note on the Frobenius-Euler numbers and polynomials associated with Bernstein polynomials. Adv. Stud. Contemp. Math. (Kyungshang) **22**(3), 399–406 (2012)
2. Araci, S., Bagdasaryan, A., Özel, C., Srivastava, H.M.: New symmetric identities involving q-zeta type functions. Appl. Math. Inf. Sci. **8**(6), 2803–2808 (2014). DOI 10.12785/amis/080616. URL http://dx.doi.org/10.12785/amis/080616
3. Bayad, A., Chikhi, J.: Apostol-Euler polynomials and asymptotics for negative binomial reciprocals. Adv. Stud. Contemp. Math. (Kyungshang) **24**(1), 33–37 (2014)
4. Bayad, A., Kim, T.: Identities involving values of Bernstein, q-Bernoulli, and q-Euler polynomials. Russ. J. Math. Phys. **18**(2), 133–143 (2011). DOI 10.1134/S1061920811020014. URL http://dx.doi.org/10.1134/S1061920811020014
5. Carlitz, L.: Degenerate Stirling, Bernoulli and Eulerian numbers. Utilitas Math. **15**, 51–88 (1979)
6. Duran, U., Acikgoz, M., Araci, S.: Symmetric identities involving weighed q-Genocchi polynomials under s_4. Proc. Jangjeon Math. **18**(4), (in press) (2015)
7. He, Y.: Symmetric identities for Carlitz's q-Bernoulli numbers and polynomials. Adv. Difference Equ. pp. 2013:246, 10 (2013). DOI 10.1186/1687-1847-2013-246. URL http://dx.doi.org/10.1186/1687-1847-2013-246
8. Kim, D.S., Kim, T.: Some identities of degenerate Euler polynomials arising from p-adic fermionic integrals on $\mathbb{Z}_p$. Integral Transforms Spec. Funct. **26**(4), 295–302 (2015). DOI 10.1080/10652469.2014.1002497. URL http://dx.doi.org/10.1080/10652469.2014.1002497
9. Kim, D.S., Lee, N., Na, J., Park, K.H.: Identities of symmetry for higher-order Euler polynomials in three variables (I). Adv. Stud. Contemp. Math. (Kyungshang) **22**(1), 51–74 (2012)
10. Kim, D.S., Lee, N., Na, J., Park, K.H.: Abundant symmetry for higher-order Bernoulli polynomials (I). Adv. Stud. Contemp. Math. (Kyungshang) **23**(3), 461–482 (2013)
11. Kim, T.: Symmetry of power sum polynomials and multivariate fermionic p-adic invariant integral on $\mathbb{Z}_p$. Russ. J. Math. Phys. **16**(1), 93–96 (2009). DOI 10.1134/S1061920809010063. URL http://dx.doi.org/10.1134/S1061920809010063
12. Kim, T.: Symmetry properties of the generalized higher-order Euler polynomials. Proc. Jangjeon Math. Soc. **13**(1), 13–16 (2010)
13. Kim, T., Dolgy, D.V., Jang, Y.S., Seo, J.J.: A note on symmetric identities for the generalized q-Euler polynomials of the second kind. Proc. Jangjeon Math. Soc. **17**(3), 375–381 (2014)

14. Kim, T., Kim, D.S., Dolgy, D.V.: Degenerate q-Euler polynomials. Adv. Difference Equ. p. 2015:246 (2015). DOI 10.1186/s13662-015-0563-y. URL http://dx.doi.org/10.1186/s13662-015-0563-y
15. Kim, Y.H., Hwang, K.W.: Symmetry of power sum and twisted Bernoulli polynomials. Adv. Stud. Contemp. Math. (Kyungshang) **18**(2), 127–133 (2009)
16. Moon, E.J., Rim, S.H., Jin, J.H., Lee, S.J.: On the symmetric properties of higher-order twisted q-Euler numbers and polynomials. Adv. Difference Equ. pp. Art. ID 765,259, 8 (2010)
17. Rim, S.H., Jeong, J.H., Lee, S.J., Moon, E.J., Jin, J.H.: On the symmetric properties for the generalized twisted Genocchi polynomials. Ars Combin. **105**, 267–272 (2012)
18. Ryoo, C.S.: A note on the weighted q-Euler numbers and polynomials. Adv. Stud. Contemp. Math. (Kyungshang) **21**(1), 47–54 (2011)
19. Şen, E.: Theorems on Apostol-Euler polynomials of higher order arising from Euler basis. Adv. Stud. Contemp. Math. (Kyungshang) **23**(2), 337–345 (2013)